



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Show that of all curves of a given length, traced on one plane between two given points, and made to revolve around a common axis situated in that plane, the Catenary generates a minimum area.

MECHANICS.

Conducted by B. F. FINKEL, Kidder, Mo. All contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

12. Proposed by J. F. W. SCHEFFER, A. M. Hagerstown, Maryland.

A horizontal table without weight is supported on three points, A , B , and C . A weight W is laid upon the table, at a point G . If $AG=a$, $BG=b$, $CG=c$, $\angle AGB=\alpha$, $\angle BGC=\beta$, and $\angle CGA=\gamma$, find the pressures upon A , B , and C .

Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

The momental equations with respect to AG , BG , and CG , are respectively:

$$P_B \times b \sin(\pi - \alpha) = P_C \times c \sin \beta. \quad \therefore \frac{P_B}{P_C} = \frac{c \sin \beta}{b \sin \alpha} \dots (m).$$

$$P_C \times c \sin(\pi - \beta) = P_A \times a \sin[\gamma - (\pi - \beta)]. \quad \therefore P_C = \left(\frac{a \sin \alpha}{c \sin \beta} \right) P_A \dots (n).$$

$$P_A \times a \sin(\pi - \gamma) = P_B \times b \sin[\alpha - (\pi - \gamma)]. \quad \therefore P_B = \left(\frac{a \sin \gamma}{b \sin \beta} \right) P_A \dots (p).$$

Put $K = \left(\frac{\sin \beta}{a} + \frac{\sin \gamma}{b} + \frac{\sin \alpha}{c} \right)$; then from the equation, $P_A + P_B + P_C = W$, we deduce the following *symmetrical* and elegant results:

$$P_A = \left(\frac{\sin \beta}{aK} \right) W, \quad P_B = \left(\frac{\sin \gamma}{bK} \right) W, \quad \text{and} \quad P_C = \left(\frac{\sin \alpha}{cK} \right) W.$$

Second Solution.

According to the logic of common-sense, why not write

$$P_A = \left(\frac{\triangle BCG}{\triangle ABC} \right) W, \quad P_B = \left(\frac{\triangle CAG}{\triangle ABC} \right) W, \quad \text{and} \quad P_C = \left(\frac{\triangle ABG}{\triangle ABC} \right) W;$$

and then with the heavy artillery of the higher mathematics *successfully defend* our position?

NOTE:—These two solutions are to take the place of the *first* solution of this problem published in the November MONTHLY.—F. P. M.